For Research Use Only

HE4 Monoclonal antibody

Catalog Number: 66557-1-Ig



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

66557-1-lg BC046106 Protein G purification GeneID (NCBI): Size: CloneNo.:

150ul, Concentration: 1500 ug/ml by 10406 5B2D9 Nanodrop and 1000 ug/ml by Bradford_{UNIPROT ID:} Recommended Dilutions:

method using BSA as the standard; Q14508 IHC 1:1000-1:5000 Source: IF/ICC 1:400-1:1600 Full Name:

Mouse WAP four-disulfide core domain 2

Isotype: Calculated MW: lgG1 13 kDa

Immunogen Catalog Number:

AG5936

Applications Tested Applications: Positive Controls:

IHC, IF/ICC, ELISA IHC: human ovary tumor tissue,

Species Specificity: IF/ICC: SKOV-3 cells,

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

WFDC2, also named as HE4, is a member of the WFDC domain family. The WFDC domain, or WAP Signature motif, contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor in many family members. HE4 is expressed in a number of normal tissues, and it is also highly expressed in a $number\ of\ tumors\ cells\ lines, such\ ovarian,\ colon,\ breast,\ lung\ and\ renal\ cells\ lines.\ In\ some\ study,\ HE4\ was\ referred$ to as a biomarker for ovarian carcinoma.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66557-1-Ig (HE4 antibody) at dilution of 1:5000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66557-1-Ig (HE4 antibody) at dilution of 1:5000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using HE4 antibody (66557-1-lg, Clone: 5B2D9) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).