

For Research Use Only

# FOXC1 Monoclonal antibody

Catalog Number: 66568-1-Ig **Featured Product**



## Basic Information

<b>Catalog Number:</b> 66568-1-Ig	<b>GenBank Accession Number:</b> NM_001453	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 150ul , Concentration: 1500 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 2296	<b>CloneNo.:</b> 1F4E11
<b>Source:</b> Mouse	<b>Full Name:</b> forkhead box C1	<b>Recommended Dilutions:</b> WB 1:2000-1:20000
<b>Isotype:</b> IgG1	<b>Calculated MW:</b> 57 kDa	
	<b>Observed MW:</b> 70-75 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : HeLa cells, HEK-293 cells, HEK293 cells, L02 cells, HepG2 cells
<b>Species Specificity:</b> Human	

## Background Information

FOXC1, also named as FKHL7 and FREAC3, binding of FREAC-3 and FREAC-4 to their cognate sites results in bending of the DNA at an angle of 80-90 degrees. Defects in FOXC1 are the cause of Axenfeld-Rieger syndrome type 3 (RIEG3). Defects in FOXC1 are the cause of iridogoniodysgenesis anomaly (IGDA). Defects in FOXC1 are a cause of Peters anomaly.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

**\*\*\* 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various cells were subjected to SDS PAGE followed by western blot with 66568-1-Ig (FOXC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of FOXC1 antibody (66568-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FOXC1 transfected HEK-293 cells.