For Research Use Only

NUP62 Monoclonal antibody

Catalog Number:66573-1-Ig




Basic Information	Catalog Number: 66573-1-lg	GenBank Accession Number BC014842	er: Purification Method: Protein G purification
	Maura		CloneNo.: 2G12B3
		^d UNIPROT ID: P37198	Recommended Dilutions: WB 1:2000-1:10000
		Full Name: nucleoporin 62kDa	IHC 1:1000-1:4000
		Calculated MW: 522 aa, 53 kDa	
		Observed MW: 62 kDa	
Applications	Tested Applications: WB. IHC. ELISA		itive Controls:
	Species Specificity: cells, PC-		B : Jurkat cells, RAW 264.7 cells, Raji cells, DU 145 ells, PC-3 cells, HepG2 cells, HSC-T6 cells
			: human prostate cancer tissue,
Background Information	NUP62 belongs to the nucleoporin NSP1/NUP62 family. It is essential component of the nuclear pore complex. The N-terminal is probably involved in nucleocytoplasmic transport. The C-terminal is probably involved in protein- protein interaction via coiled-coil formation and may function in anchorage of p62 to the pore complex. Defects in NUP62 are the cause of infantile striatonigral degeneration (SNDI). NUP62 consists of a single promoter with a CpG island and three transcribed exons. It is ubiquitously expressed, an observation compatible with its essential role in transporting cargo across the nuclear envelope. (PMID:16029492)		
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°	•	
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20°C st	0, 1	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various cells were subjected to SDS PAGE followed by western blot with 66573-1-1g (NUP62 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66573-1-1g (NUP62 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66573-1-1g (NUP62 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).