

For Research Use Only

SYK Monoclonal antibody

Catalog Number: 66721-1-Ig

Featured Product

6 Publications



Basic Information

Catalog Number: 66721-1-Ig	GenBank Accession Number: BC001645	Purification Method: Protein A purification
Size: 150ul, Concentration: 2200 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 6850	CloneNo.: 4C4A12
Source: Mouse	ENSEMBL Gene ID: ENSG00000165025	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:500-1:2000 IF-P 1:50-1:500
Isotype: IgG2b	UNIPROT ID: P43405	
Immunogen Catalog Number: AG17654	Full Name: spleen tyrosine kinase	
	Calculated MW: 635 aa, 72 kDa	
	Observed MW: 70-75 kDa	

Applications

Tested Applications: WB, IHC, IF-P, IP, ELISA	Positive Controls: WB: Raji cells, THP-1 cells, Ramos cells, Daudi cells IHC: human tonsillitis tissue, IF-P: human tonsillitis tissue,
Cited Applications: WB, IF, IP	
Species Specificity: human	
Cited Species: human, mouse, pig	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

SYK (spleen tyrosine kinase) is a key regulator of signal transduction events, apoptosis and orderly cell cycle progression in B-lineage lymphoid cells. Although SYK has not been linked to a human disease, defective expression of the closely related T-cell tyrosine kinase ZAP-70 has been associated with severe combined immunodeficiency (PMID:11494125). This protein has 2 isoforms produced by alternative splicing. SYK also promotes liver fibrosis by activation of hepatic stellate cells and acts as a biomarker for human hepatocellular carcinoma. (PMID: 29537660)

Notable Publications

Author	Pubmed ID	Journal	Application
Jie Jia	36152731	Exp Cell Res	WB
Tian-Jiao Li	35077803	Cancer Lett	WB
Xiaohua Zhang	39316625	PLoS Pathog	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66721-1-Ig (SYK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 66721-1-Ig (SYK antibody), at dilution of 1:100 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




THP-1 (WT and SYK KO) lysates prepared with RIPA buffer, 30 µg protein loaded. 66721-1-Ig incubated at 1:500 at 4°C overnight in 5% BSA in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.



WB result of SYK antibody (66721-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SYK transfected Raji cells.



MDA-MB-231 lysates prepared and IP of SYK performed using 2.0 µg of 66721-1-Ig coupled to protein G- Sepharose beads. The Ponceau stained transfers of each blot are shown. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66721-1-Ig (SYK antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).