For Research Use Only

## AFP Monoclonal antibody

Catalog Number:67852-1-lg




Basic Information	Catalog Number: 67852-1-lg	GenBank Accession Number: BC027881	Purification Method: Protein A purification
	Size: 150ul, Concentration: 1000 µg/ml by Nanodrop; Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG6089	GeneID (NCBI): 174 UNIPROT ID: P02771 Full Name: alpha-fetoprotein Calculated MW: 69 kDa Observed MW: 68-72 kDa	CloneNo.: 4B3E4 Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000
Applications	Tested Applications: WB, IHC, ELISA Species Specificity: human Note-IHC: suggested antigen ra TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	, human pl IHC : hum retrieval with rety, antigen	Controls: G2 cells, HuH-7 cells, human testis tissue, lacenta tissue nan liver tissue, human liver cancer tissue
Background Information	AFP (Alpha-fetoprotein) is a major plasma protein in the fetus and its concentration is very low in the adult (PMID:24120489). AFP can be detected at abnormally high concentrations in hepatocellular carcinomas as well as in the plasma and ascitic fluid of adults with hepatoma, indicating that AFP can serve as a tumor marker (PMID: 18669658). AFP is also a glycosylated protein and based on its binding capability to lectin Lens Culinaris Agglutinin (LCA), and total AFP can be separated into three different glycoforms, AFP-L1, AFP-L2, and AFP-L3. Core-fucosylated form of AFP (AFP-L3) is a more specific indicator than total AFP for HCC (PMID: 33128033, 35458505)		
Storage *** 20ul sizes contain 0.1% BSA	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50° Aliquoting is unnecessary for -20°C st	% glycerol pH 7.3.	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



HuH-7 cells were subjected to SDS PAGE followed by western blot with 67852-1-1g (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.





HepG2 cells were subjected to SDS PAGE followed by western blot with 67852-1-1g (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



human testis tissue were subjected to SDS PAGE followed by western blot with 67852-1-lg (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.









human placenta tissue were subjected to SDS PAGE followed by western blot with 67852-1-Ig (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67852-1-1g (AFP antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67852-1-Ig (AFP antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).





Immunohistochemical analysis of paraffinembedded human liver tissue slide using 67852-1-Ig (AFP antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded human liver tissue slide using 67852-1-Ig (AFP antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).