

Catalog Number: HZ-1038

Data Sheet



HumanKine® recombinant human VEGF165 protein

Animal Component-Free

Human cell expressed

Tag-Free


Endotoxin Free

Product Description


Animal-free Recombinant Human VEGF 165is expressed in human 293 cells as a homodimeric glycoprotein with an apparent molecular mass of 45 kDa. The E. coli expressed protein is a monomer-dimer mixture with a molecular mass 18 to 34 kDa. The cytokine is produced in a human cell expression system with serum free, chemically defined media. VEGF, also known as vascular permeability factor (VPF) or vasculotropin, is a homodimeric 34-45 kDa, heparin-binding glycoprotein with potent angiogenic, mitogenic, and vascular permeability enhancing activities specific for endothelial cells.

Alternative Names	L VEGFA, MVCD1, Vascular permeability factor, VEGF, VEGF A, VEGF165, VEGFA, VPF	
Source Human Embryonic Kidney cells (HEK293). HEK293-derived VEGF165 protein		
Species Reactivity	human	

Specifications					
Test	Method	Specification			
Activity (EC50)	Dose-dependent stimulation of the proliferation of HUVEC cells (Human Umbilical Vein Endothelial Cells)	0.3-3.75 ng/mL			
Molecular Mass	SDS-PAGE	22 to 30 and 50 kDa reduced, 40 to 50 and 75 kDa non-reduced, homodimer, glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/µg			



Recombinant human
VEGF165 (HZ-1038) induces
dose-dependent
proliferation of the HUVEC
(human umbilical vein
endothelial) cell line. Cell
number was quantitatively
assessed by PrestoBlue®
cell viability reagent. HUVEC
cells were treated with
increasing concentrations
of recombinant VEGF165 for
96 hours. The EC50 was
determined using a 4parameter non-linear



www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

	Preparation				
Shipping Temperature	ambient temperature				
Formulation 1x PBS, See Certificate of Analysis for details					
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x PBS pH 7.4 containing 0.1% endotoxin-free recombinant serum albumin (HSA). Gently swirl or tap vial to mix.				

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com