

NCOA4 Polyklonaler Antikörper

Katalog-Nr.: 10968-1-AP

1 Publikationen

Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:
10968-1-AP	BC012736
Größe:	GenID (NCBI):
150ul, Konzentration: 133 µg/ml	8031
durch die Bradford-Methode mit BSA als Standard;	Vollständiger Name:
	nuclear receptor coactivator 4
Wirt:	Berechnete Masse:
Kaninchen	70 kDa
Isotyp:	
IgG	
Immunogen Katalognummer:	
AG1408	

Anwendungen

Geprüfte Anwendungen:
ELISA
In Publikationen genannte Anwendungen:
IF, WB
Getestete Reaktivität:
Human, Maus, Ratte
Zitierte Arten:
Human

Hintergrundinformationen

Nuclear receptor coactivator 4 (NCOA4) also named androgen receptor (AR) coactivator ARA70, RFG and ELE1, is a putative co-activator that specifically enhances the activity of the androgen receptor. In human thyroid carcinomas, the Ret proto-oncogene fuses to ARA70 to form Ret/PTC3 by an intrachromosomal inversion of chromosome 10 in vivo. ARA70a can function as a ligand-enhanced co-activator of PPAR γ in adipocytes. However, PPAR γ -ARA70 transactivation can be squelched by AR, which suggests cross talk between PPAR γ - and AR-mediated response. ARA70a has no intrinsic transcription activation domain or histone acetyltransferase activity, but it interacts with histone acetyltransferase, p/CAF, CBP and p300/CBP-associated factors, and the basal transcription factor TFIIB. The interaction between ARA70 and AR occurs through the ligand-binding domain. The presence of ARA70 can enhance the androgenic activity of 17- β Estradiol (E2) and antiandrogens toward AR. ARA70 may be involved in prostate carcinogenesis and ovarian cancer and may serve as a key mediator of estrogen-androgen synergism. There are several isoforms of ZGPAT that are produced as a result of alternative splicing events. ARA70a is widely expressed, and its expression is highest in testis and adipose tissues, whereas ARA70b (The shorter variant, results from an internal 985-bp deletion.) is solely expressed in the testis. This antibody is a rabbit polyclonal antibody raised against the N-400aa of human NCOA4 protein.

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Hao Li	34917147	J Oncol	WB, IF

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten