

IMP3 Monoklonaler Antikörper

Katalog-Nr.: 66247-1-Ig

Allgemeine Informationen

Katalog-Nr.:

66247-1-Ig

Größe:

150ul, Konzentration: 3300 µg/ml von 55272

Nanodrop und 1733 µg/ml durch die Bradford-Methode mit BSA als Standard;

Wirt:

Maus

Isotyp:

IgG2b

Immunogen Katalognummer:

AG23602

GenBank-Zugangsnummer:

BC020557

GeneID (NCBI):

von 55272

Vollständiger Name:

IMP3, U3 small nucleolar ribonucleoprotein, homolog (yeast)

Berechnete Masse:

184 aa, 22 kDa

Beobachtete Masse:

22 kDa

Reinigungsmethode:

Protein-A-Reinigung

CloneNo.:

1E7D6

Empfohlene Verdünnungen:

WB 1:1000-1:4000

IHC 1:50-1:500

IF 1:200-1:800

Anwendungen

Geprüfte Anwendungen:

FC, IF, IHC, WB, ELISA

Getestete Reaktivität:

Human, Maus

Hinweis-IHC: Antigendemaskierung mit TE-Puffer pH 9,0 empfohlen. (*) Wahlweise kann die Antigendemaskierung auch mit Citratpuffer pH 6,0 erfolgen.

Positivkontrollen:

WB: HeLa-Zellen, HepG2-Zellen, SKOV-3-Zellen

IHC: humanes Mammakarzinomgewebe,

IF: HepG2-Zellen,

Hintergrundinformationen

IMP3 is a member of the insulin-like growth factor (IGF-II) mRNA binding protein (IMP) family that consist of IMP1, IMP2, and IMP3. IMP3 is an oncofetal protein associated with aggressive and advanced tumors and is specifically expressed in malignant tumors but not found in benign tissues [PMID:22497850]. IMP3 forms a ternary complex with IMP4 and MPP10 (MPHOSPH10) that interacts with U3 small nucleolar RNA (snoRNA), which is required for the early cleavage steps in pre-rRNA processing [PMID: 12655004].

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:

PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




HeLa cells were subjected to SDS PAGE followed by western blot with 66247-1-Ig (IMP3 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66247-1-Ig (IMP3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug Anti-Human IMP3 (66247-1-Ig, Clone:1E7D6) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using IMP3 antibody (66247-1-Ig, Clone: 1E7D6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using IMP3 antibody (66247-1-Ig, Clone: 1E7D6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using IMP3 antibody (66247-1-Ig, Clone: 1E7D6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).