

Nur für Forschungszwecke

Septin 9 Monoklonaler Antikörper

Katalog-Nr.:67548-1-Ig



Allgemeine Informationen

Katalog-Nr.:
67548-1-Ig

Größe:

150ul, Konzentration: 1000 µg/ml von10801

Nanodrop und 500 µg/ml durch die
Bradford-Methode mit BSA als
Standard;

Wirt:

Maus

Isotyp:

IgG1

Immunogen Katalognummer:

AG18047

GenBank-Zugangsnummer:

BC021192

GeneID (NCBI):

10801

Vollständiger Name:

septin 9

Berechnete Masse:

64 kDa

Beobachtete Masse:

42 kDa

Reinigungsmethode:

Protein-G-Reinigung

CloneNo.:

1C6F10

Empfohlene Verdünnungen:

WB 1:20000-1:100000

Anwendungen

Geprüfte Anwendungen:

WB, ELISA

Getestete Reaktivität:

Hausschwein, Human, Ratte

Positivkontrollen:

WB : Hausschwein-Kolongewebe, A549-Zellen, HeLa-Zellen, HepG2-Zellen, MCF-7-Zellen, Ratten-Kolongewebe

Hintergrundinformationen

Septin 9 belongs to the septin family which is a group of highly conserved GTP-binding proteins in eukaryotes. SEPT9 gene methylation has been implicated as a biomarker for colorectal cancer(CRC). this antibody can identify all of the isoforms about SEPT9.

Lagerung

Lagerungsbedingungen:

Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten



Various lysates were subjected to SDS PAGE followed by western blot with 67548-1-Ig (Septin 9 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 67548-1-Ig (Septin 9 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



rat colon tissue were subjected to SDS PAGE followed by western blot with 67548-1-Ig (Septin 9 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.