

Nur für Forschungszwecke

SOX1 Monoklonaler Antikörper

Katalog-Nr.:67994-1-Ig



Allgemeine Informationen

Katalog-Nr.: 67994-1-Ig	GenBank-Zugangsnummer: NM_005986	Reinigungsmethode: Protein-G-Reinigung
Größe: 150ul , Konzentration: 1000 µg/ml von6656 Nanodrop;	GeneID (NCBI): 150000	CloneNo.: 1C10H2
Wirt: Maus	Vollständiger Name: SRY (sex determining region Y)-box 1	Empfohlene Verdünnungen: WB 1:2000-1:10000
Isotyp: IgG1	Berechnete Masse: 39 kDa	
Immunogen Katalognummer: AG31118	Beobachtete Masse: 42 kDa	

Anwendungen

Geprüfte Anwendungen:
WB, ELISA

Getestete Reaktivität:
Human, Maus

Positivkontrollen:
WB : HUVEC-Zellen, ATDC-5-Zellen, COLO 320-Zellen,
HaCaT-Zellen, NCCIT-Zellen

Hintergrundinformationen

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern.

Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.


Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20ul-Größen enthalten 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Ausgewählte Validierungsdaten




HUVEC cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




ATDC-5 cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



NCCIT cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



HaCaT cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



COLO 320 cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.