

Nur für Forschungszwecke

BCAS2 Monoklonaler Antikörper

Katalog-Nr.:68003-1-Ig



Allgemeine Informationen

Katalog-Nr.: 68003-1-Ig	GenBank-Zugangsnummer: BC005285	Reinigungsmethode: Protein-G-Reinigung
Größe: 150ul , Konzentration: 1000 µg/ml von10286	GeneID (NCBI): von10286	CloneNo.: 2A12G1
Nanodrop;	Vollständiger Name: breast carcinoma amplified sequence	Empfohlene Verdünnungen: WB 1:2000-1:10000
Wirt: Maus	Berechnete Masse: 26 kDa	
Isotyp: IgG1	Beobachtete Masse: 28 kDa	
Immunogen Katalognummer: AG21309		

Anwendungen

Geprüfte Anwendungen:
WB, ELISA

Getestete Reaktivität:
Human, Maus, Ratte

Positivkontrollen:

WB : HepG2-Zellen, HEK-293-Zellen, HeLa-Zellen, Jurkat-Zellen, K-562-Zellen, LNCaP-Zellen, NIH/3T3-Zellen

Hintergrundinformationen

Breast carcinoma amplified sequence 2 (BCAS2) is preferentially known as pre-mRNA splicing factor SPF27 and was originally characterized as an up-regulated gene by amplification in human breast cancer cells. BCAS2 has been shown to be involved in DNA damage repair through the replication protein A (RPA) complex.

Lagerung

Lagerungsbedingungen:
Bei -20°C lagern. Nach dem Versand ein Jahr lang stabil

Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.

Aliquotieren ist nicht notwendig bei -20°C Lagerung

***** 20ul-Größen enthalten 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



HepG2 cells were subjected to SDS PAGE followed by western blot with 68003-1-Ig (BCAS2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



LNCaP cells were subjected to SDS PAGE followed by western blot with 68003-1-Ig (BCAS2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.