

Nur für Forschungszwecke

Phospho-MEK1 (Ser298) Monoklonaler Antikörper

Katalog-Nr.: 68047-1-Ig 1 Publikationen



Allgemeine Informationen

Katalog-Nr.:	GenBank-Zugangsnummer:	Reinigungsmethode:
68047-1-Ig	BC139729	Protein-G-Reinigung
Größe:	GenID (NCBI):	CloneNo.:
100μl, Konzentration: 1000 μg/ml von 5604 Nanodrop;	5604	3F10G10
Wirz:	Vollständiger Name:	Empfohlene Verdünnungen:
Maus	mitogen-activated protein kinase kinase 1	WB 1:5000-1:50000
Isotyp:	Berechneté Masse:	
IgG1	43 kDa	
	Beobachteté Masse:	
	40-50 kDa	

Anwendungen

Geprüfte Anwendungen:	Positivkontrollen:
WB, ELISA	WB : HeLa-Zellen, A431-Zellen, Mit Calyculin A behandelte HeLa-Zellen, Mit Nocodazol behandelte A431-Zellen
In Publikationen genannte Anwendungen:	
WB	
Getestete Reaktivität:	
Human	
Zitierte Arten:	
Maus	

Hintergrundinformationen

MAP2K1 encodes MAPK1, also known as MEK1. MEK1 variants can enhance MEK1 expression and ERK phosphorylation that together lead to continuous activation of MEK/ERK signaling pathway. MEK1 binds directly to ERK2 through a region in the N terminus of MEK1. In addition, a proline-rich (PR) regulatory sequence in MEK1 is also involved in MEK-ERK association and signal propagation. The coupling between MEK1 and ERK2 is enhanced through phosphorylation on S298 in the MEK1 PR region, whereas phosphorylation on MEK1 T292 releases the complex. MEK1 T292 is a substrate of ERK2, but the site is also phosphorylated at a basal level when ERK2 is inhibited, suggesting several regulators of this site. Although the S298 site in MEK2 has been conserved, it lacks the T292 phosphorylation site, and it is not a substrate of PAK1. (PMID: 31972311, PMID: 17928366, PMID: 22177953)

Bemerkenswerte Veröffentlichungen

Verfasser	Pubmed ID	Journal	Anwendung
Chaoqun Li	35798541	ACS Appl Mater Interfaces	WB

Lagerung


Lagerungsbedingungen:
Bei -20°C lagern.
Lagerungspuffer:
PBS mit 0.02% Natriumazid und 50% Glycerin pH 7.3.
Aliquotieren ist nicht notwendig bei -20°C Lagerung

*** 20μl-Größen enthalten 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



Non-treated A431 cells and nocodazole treated A431 cells were subjected to SDS PAGE followed by western blot with 68047-1-Ig (Phospho-MEK1 (Ser298) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.

Non-treated HeLa cells and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 68047-1-Ig (Phospho-MEK1 (Ser298) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.