

Nur für Forschungszwecke

CD8a Monoklonaler Antikörper

Katalog-Nr.: FITC-65180



Allgemeine Informationen

Katalog-Nr.:

FITC-65180

Größe:

100ug , 500 µg/ml

Wirt:

Maus

Isotyp:

IgG1, kappa

GenBank-Zugangsnummer:

BC088126

GeneID (NCBI):

24930

Vollständiger Name:

CD8a molecule

Reinigungsmethode:

Mit Protein-A-

Affinitätschromatographie gereinigt

CloneNo.:

OX-8

Anregungs-/Emissionsmaxima-

Wellenlängen:

495 nm / 524 nm

Anwendungen

Geprüfte Anwendungen:

FC

Getestete Reaktivität:

Ratte

Hintergrundinformationen

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8aβ heterodimer while CD8aα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

Lagerung

Lagerungsbedingungen:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Lagerungspuffer:

PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Ausgewählte Validierungsdaten



1X10⁶ Wistar rat splenocytes were surface stained with 0.5 ug FITC Plus Anti-Rat CD8a (FITC-65180, Clone: OX-8), or 0.5 ug Control Antibody. Cells were not fixed.