

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TSPAN15



Numéro de catalogue: 10460-1-AP

Informations de base

Numéro de catalogue:

10460-1-AP

Taille:

150ul, Concentration: 133 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG0710

Numéro d'acquisition GenBank:

BC003157

Identification du gène (NCBI):

23555

Nom complet:

tetraspanin 15

MW calculé

33 kDa

MW observés:

33 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:500-1:1000

Applications

Applications testées:

WB, ELISA

Spécificité de l'espèce:

Humain

Contrôles positifs:

WB : cancer de l'ovaire humain,

Informations générales

TSPAN15 is a membrane protein of the tetraspanin superfamily, which are characterized by the presence of four conserved transmembrane regions. Tetraspanins have been involved in diverse processes such as cell activation and proliferation, adhesion and motility, differentiation, and cancer. TSPAN15 belongs to the TSPANC8 subfamily, which also include TSPAN5, TSPAN10, TSPAN14, TSPAN17, and TSPAN33. It has been reported that TSPANC8 tetraspanins interact with ADAM10 and have a conserved function in the regulation of ADAM10 trafficking and activity, thereby positively regulating Notch receptor activation (PMID: 23091066; 23035126).

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com

W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



human ovary cancer were subjected to SDS PAGE followed by western blot with 10460-1-AP (TSPAN15 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.