

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-BTN3A2



Numéro de catalogue: 15148-1-AP

Informations de base

Numéro de catalogue: 15148-1-AP	Numéro d'acquisition GenBank: BC020214	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul, Concentration: 500 µg/ml by Nanodrop and 220 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 11118	Dilutions recommandées: WB 1:200-1:1000
Hôte: Lapin	Nom complet: butyrophilin, subfamily 3, member A2	
Isotype: IgG	MW calculé 36 kDa	
Immunogen Catalog Number: AG2212	MW observés: 36 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB: cellules SKOV-3,
Spécificité de l'espèce: Humain, souris	

Informations générales

BTN3A2, also named as Butyrophilin subfamily 3 member A2, is a 334 amino acid protein, which contains 1 Ig-like V-type (immunoglobulin-like) domain and belongs to the immunoglobulin superfamily. BTN3A2 plays a role in T-cell responses in the adaptive immune response and inhibits the release of IFNG from activated T-cells. BTN3A2 can form Homodimer and is detected in T-cells and natural killer cells.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



SKOV-3 cells were subjected to SDS PAGE followed by western blot with 15148-1-AP (BTN3A2 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



SKOV-3 cells were subjected to SDS PAGE followed by western blot with 15148-1-AP (BTN3A2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.