

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TFAP2D



Numéro de catalogue: 18835-1-AP

Informations de base

Numéro de catalogue:
18835-1-AP

Taille:
150ul, Concentration: 800 µg/ml by
Nanodrop and 367 µg/ml by Bradford
method using BSA as the standard;

Hôte:
Lapin

Isotype:
IgG

Numéro d'acquisition GenBank:
BC166653

Identification du gène (NCBI):
83741

Nom complet:
transcription factor AP-2 delta
(activating enhancer binding protein
2 delta)

MW calculé
50 kDa

MW observés:
47-52 kDa

Méthode de purification:
Purification par affinité contre
l'antigène

Dilutions recommandées:
WB 1:500-1:1000

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, rat, souris

Contrôles positifs:

WB: cellules HeLa,

Informations générales

TFAP2D, also named as TFAP2BL1, belongs to the AP-2 family. TFAP2D is a sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC. It is highly expressed in adult thymus, prostate, small intestine, skeletal muscle, placenta, brain, and testis tissues.

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azote de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



HeLa cells were subjected to SDS PAGE followed by western blot with 18835-1-AP (TFAP2D Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.