

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-FIZ1



Numéro de catalogue: 23257-1-AP

Informations de base

Numéro de catalogue:

23257-1-AP

Taille:

150ul, Concentration: 587 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG19760

Numéro d'acquisition GenBank:

BC136512

Identification du gène (NCBI):

84922

Nom complet:

FLT3-interacting zinc finger 1

MW calculé

496 aa, 52 kDa

MW observés:

52 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:200-1:1000

Applications

Applications testées:

WB, ELISA

Spécificité de l'espèce:

Humain, souris

Contrôles positifs:

WB : cellules Jurkat, cellules HEK-293, cellules NIH/3T3

Informations générales

FLT3 interacting zinc finger 1 (FIZ1) is a ubiquitous expressed protein that composes of 11 C2H2 zinc finger domains spread over the entire protein in clusters of two, three, or four fingers. The interaction of FIZ1 was mapped to the catalytic domain of FLT3, which is a receptor tyrosine kinase involved in proliferation and survival of hematopoietic progenitor cells as well as differentiation of early B lymphoid progenitors (10409713). In the mammalian retina, FIZ1 interacts with NRL (Neural-Retina Leucine-zipper), an essential transcriptional activator of rod photoreceptor-specific genes, and the concentration of FIZ1 in the retina will increase during photoreceptor terminal maturation (18854042).

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Jurkat cells were subjected to SDS PAGE followed by western blot with 23257-1-AP (FIZ1 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.