

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-SLC25A22



Numéro de catalogue: 25402-1-AP

1 Publications

Informations de base

Numéro de catalogue:

25402-1-AP

Taille:

150ul, Concentration: 1000 µg/ml by Nanodrop and 460 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG19958

Numéro d'acquisition GenBank:

BC023545

Identification du gène (NCBI):

79751

Nom complet:

solute carrier family 25 (mitochondrial carrier: glutamate), member 22

MW calculé

323 aa, 34 kDa

MW observés:

30-34 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:1000-1:6000

IHC 1:50-1:500

IF 1:50-1:500

Applications

Applications testées:

IF, IHC, WB, ELISA

Demandes citées:

WB

Spécificité de l'espèce:

Humain, rat, souris

Espèces citées:

souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) A défaut, 'le démasquage de l'antigène peut être 'effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : tissu cérébral de souris, cellules Caco-2, cellules HT-29, cellules NCI-H1299, tissu cérébral de rat

IHC : tissu cérébral de souris, tissu de cancer du côlon humain, tissu de cancer du foie humain

IF : cellules COS-7, tissu cérébral de souris

Informations générales

SLC25A22 encodes a mitochondrial glutamate/H⁺ symporter with highly expression in brain, especially in astrocytes, where it controls glutamate uptake. The mutations of SLC25A22 are associated with the early/infantile epileptic encephalopathies. It has been shown that SLC25A22 plays a key role in promoting proliferation and migration in cancer, such as human colorectal cancer (CRC) and Gallbladder cancer (GBC). And there is a result that the expression of SLC25A22 in GBC was significantly higher than that in adjacent tissues. 25402-1-AP antibody is specific to SLC25A22.(PMID: 30814911, 25033742, 24596948)

Publications notables

Autrice	Pubmed ID	Journal	Application
Maria F Noterman	33836587	Proc Natl Acad Sci U S A	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Immunofluorescent analysis of (4% PFA) fixed COS-7 cells using 25402-1-AP (SLC25A22 antibody), at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 25402-1-AP (SLC25A22 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 25402-1-AP (SLC25A22 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 25402-1-AP (SLC25A22 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.