

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-BOK



Numéro de catalogue: 26430-1-AP

Informations de base

Numéro de catalogue: 26430-1-AP	Numéro d'acquisition GenBank: BC006203	Méthode de purification: Purification par affinité contre l'antigène
Taille: 150ul , Concentration: 300 µg/ml by Nanodrop;	Identification du gène (NCBI): 666	Dilutions recommandées: WB 1:500-1:2000
Hôte: Lapin	Nom complet: BCL2-related ovarian killer	
Isotype: IgG	MW calculé 23 kDa	
Immunogen Catalog Number: AG24070	MW observés: 23 kDa	

Applications

Applications testées: FC, WB, ELISA	Contrôles positifs: WB : cellules HepG2,
Spécificité de l'espèce: Humain	

Informations générales

BOK belongs to the BCL2 family, members of which form homo- or heterodimers, and act as anti- or proapoptotic regulators that are involved in a wide variety of cellular processes. Studies in rat show that this protein has restricted expression in reproductive tissues, interacts strongly with some antiapoptotic BCL2 proteins, not at all with proapoptotic BCL2 proteins, and induces apoptosis in transfected cells. Thus, this protein represents a proapoptotic member of the BCL2 family.

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



HepG2 cells were subjected to SDS PAGE followed by western blot with 26430-1-AP (BOK antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1×10^6 MCF-7 cells were intracellularly stained with 0.4 μ g Anti-Human BOK (26430-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 μ g Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).