

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-TMIGD1



Numéro de catalogue: 27174-1-AP

2 Publications

## Informations de base

Numéro de catalogue:

27174-1-AP

Taille:

150ul, Concentration: 260 µg/ml by Nanodrop and 233 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG25691

Numéro d'acquisition GenBank:

BC137201

Identification du gène (NCBI):

388364

Nom complet:

transmembrane and immunoglobulin domain containing 1

MW calculé

262 aa, 29 kDa

MW observés:

45 kDa, 29 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:200-1:1000

IHC 1:50-1:500

## Applications

Applications testées:

IHC, WB, ELISA

Demandes citées:

IF, IHC, WB

Spécificité de l'espèce:

Humain, souris

Espèces citées:

Humain, souris

Contrôles positifs:

WB : tissu rénal de souris,

IHC : tissu rénal de souris, tissu de côlon humain

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) A défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.**

## Informations générales

TMIGD1 (Transmembrane and immunoglobulin domain containing 1) is involved in various cellular processes such as regulation of cell structure, cell-cell adhesion, and cellular movement. It protects cells from oxidative- and nutrient-deprivation-induced cell injury by stabilizing the transepithelial electric resistance and permeability of renal epithelial cells. TMIGD1 has a calculated molecular mass of 29 kDa, and can be detected as 45 kDa due to the N-glycosylation (PMID: 26342724).

## Publications notables

| Autrice            | Pubmed ID | Journal             | Application |
|--------------------|-----------|---------------------|-------------|
| Christian Hartmann | 36099341  | Sci Signal          | WB, IF      |
| Xin Chang          | 37301121  | Int Immunopharmacol | WB, IHC     |

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20°C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 27174-1-AP (TMIGD1 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 27174-1-AP (TMIGD1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 27174-1-AP (TMIGD1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).