

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-TRIM3



Numéro de catalogue: 28392-1-AP

## Informations de base

Numéro de catalogue:

28392-1-AP

Taille:

150ul, Concentration: 700 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG28823

Numéro d'acquisition GenBank:

NM\_001248006

Identification du gène (NCBI):

10612

Nom complet:

tripartite motif-containing 3

MW calculé

81 kDa

MW observés:

75 kDa, 80-85 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:1000-1:6000

## Applications

Applications testées:

WB, ELISA

Spécificité de l'espèce:

Humain, rat, souris

Contrôles positifs:

WB : tissu cérébral de souris, tissu cérébral de rat

## Informations générales

TRIM3, a member of the tripartite motif (TRIM) family, also called the 'RING-B-box-coiled-coil' (RBCC) subgroup of RING finger proteins. The TRIM family of genes is largely studied because of their roles in development, differentiation and host cell antiviral defenses. TRIM3 is a mammalian tumor suppressor whose loss can increase the incidence and promote the development of GBM (PMID: 23318451). TRIM3 has four isoforms with the MW of 67-80 kDa and can be detected as 75, 80, 82, 85 kDa after posttranslational modification (PMID: 24947043/PMID: 24086586).

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 28392-1-AP (TRIM3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.