

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti- Phospho-p70(S6K) (Thr229)

Numéro de catalogue: 28988-1-AP



## Informations de base

Numéro de catalogue:	BC053365	Méthode de purification:
28988-1-AP		Purification par affinité contre l'antigène
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
100ul , Concentration: 800 µg/ml by Nanodrop;	6198	WB 1:1000-1:6000
Hôte:	Nom complet:	
Lapin	ribosomal protein S6 kinase, 70kDa, polypeptide 1	
Isotype:	MW calculé	
IgG	59 kDa	
	MW observés:	
	65-85 kDa	

## Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : cellules HeLa traitées à la λ phosphatase,
Spécificité de l'espèce:	
Humain	

## Informations générales

The Rps6kb1 gene encodes the 70 kDa ribosomal protein S6 kinase (p70S6K), which is a serine/threonine kinase regulated by phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin (mTOR) pathway. P70S6K plays a crucial role in controlling cell cycle, growth and survival. The PI3K/mTOR signalling pathway is one of the major mechanisms for controlling cell survival, proliferation and metabolism and is the central regulator of translation of some components of protein synthesis system. Due to alternative translation two isoform S6K1 proteins are known to exist in mammalian cells: p85 S6K1 and p70 S6K1, which is identical to p85 S6K but lacks its first 23 amino acids. In addition, mammalian cells express a second S6K1 isoform spanning 316 amino acids (p31 S6K1). (PMID: 25100792, PMID: 24970012, PMID: 21602892)

## Stockage


Stockage:  
Stocker à -20 °C  
Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Non-treated HeLa, phosphatase inhibitor treated and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 28988-1-AP (Phospho-p70(S6K) (Thr229) antibody) at dilution of 1:3000 incubated at room temperature for 1 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.