

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-HADHB

Numéro de catalogue: 29091-1-AP

2 Publications



Informations de base

Numéro de catalogue:	BC017564	Méthode de purification:
29091-1-AP		Purification par affinité contre l'antigène
Taille:	3032	Dilutions recommandées:
150ul , Concentration: 350 µg/ml by Nanodrop;		WB 1:1000-1:8000
Hôte:	hydroxyacyl-Coenzyme A dehydrogenase/3-ketoacyl-Coenzyme A thiolase/enoyl-Coenzyme A hydratase (trifunctional protein), beta subunit	
Lapin		
Isotype:		
IgG		
Immunogen Catalog Number:	51 kDa	
AG30298		
Nom complet:	52 kDa	
MW calculé		
MW observés:		

Applications

Applications testées:	WB, ELISA	Contrôles positifs:
		WB : cellules HepG2, cellules MCF-7, tissu cardiaque de rat, tissu cardiaque de souris, tissu de muscle squelettique de souris, tissu hépatique de rat
Demandes citées:	IHC, WB	
Spécificité de l'espèce:	Humain, rat, souris	
Espèces citées:	Humain	

Informations générales

HADHB, also named as TP- beta, Acetyl-CoA acyltransferase and Beta-ketothiolase, is a mitochondrial trifunctional enzyme subunit beta. Mitochondrial trifunctional enzyme catalyzes the last three of the four reactions of the mitochondrial beta-oxidation pathway. The mitochondrial beta-oxidation pathway is the major energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA. Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids. Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities, while the trifunctional enzyme subunit beta/HADHB described here bears the 3-ketoacyl-CoA thiolase activity. HADHB has 2 isoforms produced by alternative splicing with the MW of 49 kDa and 51 kDa.

Publications notables

Autrice	Pubmed ID	Journal	Application
Yun Li	36518312	Front Oncol	WB,IHC
Olga Gourdomichali	35205152	Biology (Basel)	WB

Stockage

Stockage:
Stocker à -20 °C.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 29091-1-AP (HADHB antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 29091-1-AP (HADHB antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).