

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-TRA2B



Numéro de catalogue: 29627-1-AP

1 Publications

## Informations de base

<b>Numéro de catalogue:</b> 29627-1-AP	<b>Numéro d'acquisition GenBank:</b> BC005898	<b>Méthode de purification:</b> Purification par affinité contre l'antigène
<b>Taille:</b> 150ul , Concentration: 850 µg/ml by Nanodrop;	<b>Identification du gène (NCBI):</b> 6434	<b>Dilutions recommandées:</b> WB 1:1000-1:4000
<b>Hôte:</b> Lapin	<b>Nom complet:</b> transformer 2 beta homolog (Drosophila)	
<b>Isotype:</b> IgG	<b>MW calculé:</b> 34 kDa	
<b>Immunogen Catalog Number:</b> AG30637	<b>MW observés:</b> 34 kDa	

## Applications

<b>Applications testées:</b> WB, ELISA	<b>Contrôles positifs:</b> WB : cellules HEK-293, cellules HeLa, cellules Jurkat, cellules NIH/3T3
<b>Demandes citées:</b> IHC, WB	
<b>Spécificité de l'espèce:</b> Humain, souris	
<b>Espèces citées:</b> Humain	

## Informations générales

Transformer-2 protein homolog beta, also known as TRA2B, is a sequence-specific RNA-binding protein that participates in the control of pre-mRNA splicing. TRA2B can alter pre-mRNA splicing patterns by antagonizing the effects of splicing regulators, like RBMX.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Mo Juan Li	36861193	Biomed Environ Sci	WB,IHC

## Stockage


**Stockage:**  
Stocker à -20 °C.  
**Tampon de stockage:**  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



HEK-293 cells were subjected to SDS PAGE followed by western blot with 29627-1-AP (TRA2B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 29627-1-AP (TRA2B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.