

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti- Phospho-PLC_{G1} (Ser1248)

Numéro de catalogue: 29676-1-AP

1 Publications



Informations de base

Numéro de catalogue:	BC144136	Méthode de purification:
29676-1-AP	Purification par affinité contre l'antigène	
Taille:	Identification du gène (NCBI):	Dilutions recommandées:
100ul , Concentration: 300 µg/ml by Nanodrop;	5335	WB 1:500-1:2000
Hôte:	Nom complet:	
Lapin	phospholipase C, gamma 1	
Isotype:	MW observés:	
IgG	145 kDa	

Applications

Applications testées:	Contrôles positifs:
WB, ELISA	WB : cellules Jurkat traitées à la calyculine A,
Demandes citées:	
WB	
Spécificité de l'espèce:	
Humain	
Espèces citées:	
Humain	

Informations générales

Phosphoinositide phospholipase C-gamma-1 (PLC_{G1}) which belongs to the phosphoinositide-specific phospholipase C (PLC) family, is activated by many extracellular stimuli including hormones, neurotransmitters, and growth factors and modulates several cellular and physiological functions necessary for tumorigenesis such as cell survival, migration, invasion, and angiogenesis by generating inositol 1,4,5-triphosphate (IP₃) and diacylglycerol (DAG) via hydrolysis of phosphatidylinositol 4,5-bisphosphate (PIP₂) (PMID: 9242915). Phosphorylation is one of the key mechanisms that regulate the activity of PLC. PLC_γ is activated by both receptor and non-receptor tyrosine kinases (PMID: 2472218). PLC_γ forms a complex with EGF and PDGF receptors, which leads to the phosphorylation of PLC_γ at Tyr771, 783, and 1248 (PMID: 1708307). It has also been shown that PKA-mediated phosphorylation at Ser1248 is inhibitory to PLC_γ 1 tyrosine phosphorylation and phospholipase activity in CD3-treated Jurkat cells (PMID: 1370476), suggesting that Ser1248 may be an allosteric regulator of PLC_γ 1 activity.

Publications notables

Autrice	Pubmed ID	Journal	Application
Yao-Qi Zhou	37012514	Cell Oncol (Dordr)	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20°C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Non-treated and Calyculin A treated Jurkat cells were subjected to SDS PAGE followed by western blot with 29676-1-AP (Phospho-PLCG1 (Ser1248) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.