

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-TSPAN13



Numéro de catalogue: 30072-1-AP

Informations de base

Numéro de catalogue:

30072-1-AP

Taille:

150ul, Concentration: 600 µg/ml by Nanodrop;

Hôte:

Lapin

Isotype:

IgG

Immunogen Catalog Number:

AG32345

Numéro d'acquisition GenBank:

BC033863

Identification du gène (NCBI):

27075

Nom complet:

tetraspanin 13

MW calculé

204 aa, 22 kDa

MW observés:

28-35 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:2000-1:16000

Applications

Applications testées:

WB, ELISA

Spécificité de l'espèce:

Humain

Contrôles positifs:

WB : cellules PC-3,

Informations générales

TSPAN13 (also known as NET6 or TM4SF13) is a membrane protein of the tetraspanin superfamily, which are characterized by the presence of four conserved transmembrane regions. Many of these members have been implicated in signal transduction, cell-cell interactions, and cellular activation and development. Overexpression of TSPAN13 has been reported in prostate cancer, suggesting that TSPAN13 may have an important role in the progression of prostate cancer (PMID: 19148481). The experimentally determined molecular mass of 28-35 kDa is larger than the calcular molecular weight of 22 kDa, which could be a result of post-translational modifications (PMID: 23648579).

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquoteage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com

W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



PC-3 cells were subjected to SDS PAGE followed by western blot with 30072-1-AP (TSPAN13 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.