

À des fins de recherche uniquement

Anticorps Polyclonal de lapin anti-HIPK2



Numéro de catalogue: 55408-1-AP

Phare

5 Publications

Informations de base

Numéro de catalogue:

55408-1-AP

Taille:

150ul, Concentration: 1000 µg/ml by Nanodrop and 307 µg/ml by Bradford method using BSA as the standard;

Hôte:

Lapin

Isotype:

IgG

Numéro d'acquisition GenBank:

NM_022740

Identification du gène (NCBI):

28996

Nom complet:

homeodomain interacting protein kinase 2

MW calculé

131 kDa

MW observés:

131 kDa, 101 kDa

Méthode de purification:

Purification par affinité contre l'antigène

Dilutions recommandées:

WB 1:200-1:1000

Applications

Applications testées:

WB, ELISA

Demandes citées:

IF, IHC, WB

Spécificité de l'espèce:

Humain, souris

Espèces citées:

Humain, souris

Contrôles positifs:

WB : tissu rénal de souris,

Informations générales

HIPK2 is a serine/threonine kinase participating in transcriptional regulation, growth control, and apoptosis. The kinase contributes to the regulation of remarkably diverse pathways such as p53 activation or Wnt signaling. There is very low levels of HIPK2 protein expression in several thyroid tumors and the combination of HIPK2 and Gal-3, would be helpful in ameliorating the preoperative recognition of thyroid cancer. HIPK2 may represent a new tumour suppressor gene for these types of cancers and may constitute a new potential promising diagnostic marker and therapeutic target. (PMID:21698151).

Publications notables

Autrice	Pubmed ID	Journal	Application
Qiulian Zhou	36182775	EBioMedicine	IF
Hui-Ying Hu	30936013	Int J Biol Macromol	WB
Qiqi Yu	34973299	Microvasc Res	IHC

Stockage

Stockage:

Stocker à -20 °C.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 55408-1-AP (HIPK2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.