

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD8a

Numéro de catalogue: 65205-1-Ig



## Informations de base

Numéro de catalogue:

65205-1-Ig

Taille:

100ug, 0.5 mg/ml

Hôte:

Rat

Isotype:

IgG2b, lambda

Numéro d'acquisition GenBank:

BC030679

Identification du gène (NCBI):

12525

Nom complet:

CD8 antigen, alpha chain

Méthode de purification:

Purification par protéine A

CloneNo.:

5H10-1

## Applications

Applications testées:

FC

Spécificité de l'espèce:

souris

## Informations générales

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8aβ heterodimer while CD8aa homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).

## Stockage

Stockage:

Store at 2-8°C. Stable for one year after shipment.

Tampon de stockage:

PBS avec azoture de sodium à 0,1 %, pH 7,3.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> C57BL/c mouse splenocytes were surface co-stained with APC Anti-Mouse CD3ε and 0.25 μg Anti-Mouse CD8a (65205-1-Ig, Clone:5H10-1) and FITC anti-rat IgG2b Antibody at dilution 1:100, or rat IgG2b Isotype Control and FITC anti-rat IgG2b Antibody at dilution 1:100. Cells were not fixed.