

À des fins de recherche uniquement

Anticorps Monoclonal anti-TRIP13

Numéro de catalogue: 67759-1-Ig

Phare

1 Publications



Informations de base

Numéro de catalogue: 67759-1-Ig	Numéro d'acquisition GenBank: BC000404	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 9319	CloneNo.: 1F1E9
Hôte: Mouse	Nom complet: thyroid hormone receptor interactor 13	Dilutions recommandées: WB 1:2000-1:10000
Isotype: IgG1	MW calculé 49 kDa	
Immunogen Catalog Number: AG6595	MW observés: 49 kDa	

Applications

Applications testées:

WB, ELISA

Demandes citées:

WB

Spécificité de l'espèce:

Humain, rat, souris

Espèces citées:

Humain

Contrôles positifs:

WB : cellules HEK-293, cellules 4T1, cellules HeLa, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules LNCaP, cellules NIH/3T3, cellules U2OS

Informations générales

TRIP13 gene encodes thyroid receptor-interacting protein 13, that interact with thyroid hormone receptors, also known as human papillomavirus type 16 E1 protein-binding protein (16E1BP). This protein belongs to the AAA ATPase family and the PCH2 subfamily. TRIP13 gene may be one of the several playing roles in early-stage non-small cell lung cancer. TRIP13 also functions in chromosome recombination and chromosome structure development during meiosis, and it is required for development of higher-order chromosome structures and is needed for synaptonemal-complex formation.

Publications notables

Autrice	Pubmed ID	Journal	Application
Xiaoying Liu	34061428	Environ Toxicol	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67759-1-Ig (TRIP13 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.