

À des fins de recherche uniquement

Anticorps Monoclonal anti-SOX1

Numéro de catalogue: 67994-1-Ig



Informations de base

Número de catalogue: 67994-1-Ig	Número d'acquisition GenBank: NM_005986	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 6656	CloneNo.: 1C10H2
Hôte: Mouse	Nom complet: SRY (sex determining region Y)-box 1	Dilutions recommandées: WB 1:2000-1:10000
Isotype: IgG1	MW calculé 39 kDa	
Immunogen Catalog Number: AG31118	MW observés: 42 kDa	

Applications

Applications testées:
WB, ELISA
Spécificité de l'espèce:
Humain, souris

Contrôles positifs:

WB : cellules HUVEC, cellules ATDC-5, cellules COLO
320, cellules HaCaT, cellules NCCIT

Informations générales

Stockage

Stockage:
Stocker à -20 °C
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech
Group brand and is not available to purchase from any
other manufacturer.


Données de validation sélectionnées




HUVEC cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




ATDC-5 cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



NCCIT cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



HaCaT cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



COLO 320 cells were subjected to SDS PAGE followed by western blot with 67994-1-Ig (SOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.