

À des fins de recherche uniquement

# Anticorps Monoclonal anti-MYBBP1A

Numéro de catalogue: 67996-1-Ig



## Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
67996-1-Ig	BC050546	Purification par protéine G
Taille:	Identification du gène (NCBI):	CloneNo.:
150ul , Concentration: 1000 µg/ml by Nanodrop;	10514	1H2D10
Hôte:	Nom complet:	Dilutions recommandées:
Mouse	MYB binding protein (P160) 1a	WB 1:5000-1:50000
Isotype:	MW calculé	IHC 1:500-1:2000
IgG1	149 kDa	IF 1:300-1:1200
Immunogen Catalog Number:	MW observés:	
AG6008	140 kDa	

## Applications

Applications testées:	Contrôles positifs:
IF, IHC, WB, ELISA	WB : cellules LNCaP, cellules 4T1, cellules HEK-293, cellules HeLa, cellules HepG2, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules NIH/3T3
Spécificité de l'espèce:	IHC : tissu rénal humain,
Humain, rat, souris	IF : cellules HepG2, cellules NIH/3T3

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (\*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6.0.**

## Informations générales

The protooncogene MYB is predominantly expressed in immature hemopoietic cells where it has an essential role in hemopoietic cell proliferation and differentiation. Oncogenically activated forms of MYB is generally N- and/or C-terminal truncations of the normal MYB protein. Removal of the C terminus of MYB disrupts or deletes a region termed the negative regulatory domain (NRD), resulting in an increase in DNA binding, transactivation, and transformation by MYB. One feature of the NRD is a leucine zipper-like motif [PMID: 8302594]. Murine Myb-binding protein-1a (MYBBP1A), originally called P160, was identified by its ability to interact specifically with Myb via this leucine zipper-like motif. MYBBP1A modulates MYB activity upon binding to the MYB NRD [PMID: 10644447, 9447996].

## Stockage

Stockage:  
Stocker à -20 °C.  
Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20°C


\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.






## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67996-1-Ig (MYBBP1A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 67996-1-Ig (MYBBP1A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MYBBP1A antibody (67996-1-Ig, Clone: 1H2D10) at dilution of 1:600 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).