

À des fins de recherche uniquement

Anticorps Monoclonal anti-BCAS2

Numéro de catalogue: 68003-1-Ig



Informations de base

Numéro de catalogue: 68003-1-Ig	Numéro d'acquisition GenBank: BC005285	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 10286	CloneNo.: 2A12G1
Hôte: Mouse	Nom complet: breast carcinoma amplified sequence WB 1:2000-1:10000 2	Dilutions recommandées:
Isotype: IgG1	MW calculé 26 kDa	
Immunogen Catalog Number: AG21309	MW observés: 28 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, rat, souris

Contrôles positifs:

WB : cellules HepG2, cellules HEK-293, cellules HeLa,
cellules HSC-T6, cellules Jurkat, cellules K-562,
cellules LNCaP, cellules NIH/3T3

Informations générales

Breast carcinoma amplified sequence 2 (BCAS2) is preferentially known as pre-mRNA splicing factor SPF27 and was originally characterized as an up-regulated gene by amplification in human breast cancer cells. BCAS2 has been shown to be involved in DNA damage repair through the replication protein A (RPA) complex.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



HepG2 cells were subjected to SDS PAGE followed by western blot with 68003-1-Ig (BCAS2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



LNCaP cells were subjected to SDS PAGE followed by western blot with 68003-1-Ig (BCAS2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.