

À des fins de recherche uniquement

# Anticorps Monoclonal anti-IGF2BP3

Numéro de catalogue: 68470-1-Ig



## Informations de base

Numéro de catalogue: 68470-1-Ig	Numéro d'acquisition GenBank: BC065269	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 10643	CloneNo.: 1G6A7
Hôte: Mouse	Nom complet: insulin-like growth factor 2 mRNA binding protein 3	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 64 kDa	
Immunogen Catalog Number: AG6349	MW observés: 64 kDa	

## Applications

Applications testées:  
WB, ELISA

Spécificité de l'espèce:  
Humain, rat, souris

Contrôles positifs:

WB : cellules HeLa, cellules A549, cellules HEK-293,  
cellules HepG2, cellules HSC-T6, cellules Jurkat,  
cellules K-562, cellules NIH/3T3

## Informations générales

IGF2BP3, also named as IMP3, KOC1 and VICKZ3, belongs to the RRM IMP/VICKZ family. It is one of the RNA binding proteins involved in mRNA localization and translational control. IGF2BP3 is expressed during embryogenesis, as well as in some malignant tumors. It can be used as an independent prognostic factor for osteosarcoma.

## Stockage

Stockage:  
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



A549 cells were subjected to SDS PAGE followed by western blot with 68470-1-Ig (IGF2BP3 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68470-1-Ig (IGF2BP3 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.