

À des fins de recherche uniquement

Anticorps Monoclonal anti-TIA1

Numéro de catalogue: 68486-1-Ig



Informations de base

Numéro de catalogue: 68486-1-Ig	Numéro d'acquisition GenBank: BC015944	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 7072	CloneNo.: 2C11F4
Hôte: Mouse	Nom complet: TIA1 cytotoxic granule-associated RNA binding protein	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 214 aa, 24 kDa, 43 kDa	
Immunogen Catalog Number: AG2778	MW observés: 38-40 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, rat, souris

Contrôles positifs:

WB : cellules HUVEC, cellules COLO 320, cellules HepG2, cellules HSC-T6, cellules hTERT-RPE1, cellules Jurkat, cellules K-562, cellules MOLT-4, cellules NIH/3T3

Informations générales

TIA1, also named as p40-TIA-1, is involved in alternative pre-RNA splicing and regulation of mRNA translation by binding to AU-rich elements (AREs) located in mRNA 3' untranslated regions (3' UTRs). It possesses nucleolytic activity against cytotoxic lymphocyte target cells. TIA1 may be involved in apoptosis. Two isoforms of this protein exist - 41kDa and 42kDa. one of these was a missense variant (P362L) in TIA1. Similar to the ALS-related disease proteins TDP-43, hnRNPA1, and FUS, TIA1 is an RNA-binding protein containing a prionlike LCD and assembles into membrane-less organelles, including SGs. Postmortem neuropathology of five TIA1 mutations carriers showed a consistent pathological signature with numerous round, hyaline, TAR DNA-binding protein 43 (TDP-43)-positive inclusions. TIA1 mutations significantly increased the propensity of TIA1 protein to undergo phase transition. In live cells, TIA1 mutations delayed stress granule (SG) disassembly and promoted the accumulation of non-dynamic SGs that harbored TDP-43.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68486-1-Ig (TIA1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.