

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD8a

Numéro de catalogue:**CL405-65205**



## Informations de base

Numéro de catalogue:	BC030679	Méthode de purification:
CL405-65205		Purification par affinité
Taille:	Identification du gène (NCBI):	CloneNo.:
100ug , 0.5 mg/ml	12525	5H10-1
Hôte:	Nom complet:	Excitation/Emission maxima
Rat	CD8 antigen, alpha chain	wavelengths:
Isotype:		399 nm / 422 nm
IgG2b, lambda		

## Applications

Applications testées:  
FC

Spécificité de l'espèce:  
souris

## Informations générales

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer or CD8α or as a heterodimer of CD8α and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αα homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).


## Stockage

Stockage:  
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.  
Tampon de stockage:  
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> mouse splenocytes were surface co-stained with CoraLite® Plus 647 Anti-Mouse CD3 and 0.5 ug CoraLite® Plus 405 Anti-Mouse CD8a (CL405-65205, Clone:5H10-1) or 0.5 ug Control Antibody. Cells were not fixed.