

À des fins de recherche uniquement

# Anticorps Polyclonal de lapin anti-ARF4

Numéro de catalogue:**CL647-11673** Phare



## Informations de base

|  |                                |   |
|--|--------------------------------|---|
| Numéro de catalogue:                           | BC003364                       | Méthode de purification:                    |
| CL647-11673                                    |                                | Purification par affinité contre l'antigène |
| Taille:  | Identification du gène (NCBI): | Excitation/Emission maxima wavelengths:     |
| 100ul , Concentration: 1000 µg/ml by Nanodrop; | 378                            | 654 nm / 674 nm                             |
| Hôte:  | Nom complet:                   |   |
| Lapin  | ADP-ribosylation factor 4      |   |
| Isotype:                                       | MW calculé                     |   |
| IgG  | 180 aa, 21 kDa                 |   |
| Immunogen Catalog Number:                      | MW observés:                   |   |
| AG2272   | 21 kDa                         |   |

## Applications

### Applications testées:

FC (Intra)

### Spécificité de l'espèce:

Humain, rat, souris

## Informations générales

ARF4, also named as ARF2, belongs to the small GTPase superfamily and Arf family. ADP-ribosylation factors (ARFs) are members of the ARF family of GTP-binding proteins of the Ras superfamily, with 20 kDa protein size. ARFs bind and regulate GTP/GDP cycle by alternating between the active GTP-bound and inactive GDP-bound conformations. ARF family proteins are essential and ubiquitous in eukaryotes. Six highly conserved members of the family have been identified in mammalian cells. They function in vesicular traffic and actin remodelling and other bioprocesses in cells. ARF4 and ARF5 are class II ARFs. Their function are not fully described. (PMID: 7759471, PMID: 16042562). This antibody can bind ARFs for the close sequences.

## Stockage

### Stockage:

Stocker à -20 °C. Éviter toute exposition à la lumière. Stable pendant un an après l'expédition.

### Tampon de stockage:

PBS avec glycérol à 50 %, Proclin300 à 0,05 % et BSA à 0,5 %, pH 7,3.


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



$1 \times 10^6$  HepG2 cells were intracellularly stained with 0.2 ug CoralLite® Plus 647 Anti-Human ARF4 (CL647-11673) (red), or 0.2 ug isotype control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).