

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD279 (PD-1)



Numéro de catalogue:[CL647-65142](#)

Informations de base

Numéro de catalogue:	BC119179	Méthode de purification:
CL647-65142	Purification par affinité	
Taille:	Identification du gène (NCBI):	
100ug , 500 µg/ml	18566	
Hôte:	Nom complet:	
Rat	programmed cell death 1	
Isotype:		Excitation/Emission maxima
IgG2b, kappa		wavelengths:
		654 nm / 674 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

souris

Informations générales

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage:

PBS avec azoture de sodium à 0,1 % et BSA à 0,5 %, pH 7,3.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ Con-A stimulated BALB/c mouse
splenocytes were surface stained with 0.5 ug
CoraLite® Plus 647 Anti-Mouse CD279 (PD-1)
(CL647-65142, Clone: RMP1-30) or isotype control
antibody. Cells were not fixed.