## À des fins de recherche uniquement

## Anticorps Monoclonal anti-CD2

Numéro de catalogue: CR-65192



Informations de base

Numéro de catalogue:

100 tests , 5  $\mu$ l/test

Numéro d'acquisition GenBank: BC033583

Méthode de purification: Purification par affinité

CR-65192

Identification du gène (NCBI):

CloneNo.:

Taille:

TS1/8

Hôte: Mouse Nom complet: CD2 molecule MW calculé

Excitation/Emission maxima wavelengths:

Isotype: IgG1, kappa

351 aa, 39 kDa

592 nm / 611 nm

**Applications** 

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD2 is a cell surface glycoprotein present on a majority of thymocytes, all mature T cells and subset of NK cells but not on B lymphocytes. It is a pan T-cell marker. CD2 interacts with lymphocyte function-associated antigen (LFA-3/CD58) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

Stockage


Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage

PBS with 0.09% sodium azide and 0.5% BSA.

## Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with CL647 Anti-Human CD3 and 5 ul Cardinal Red™ Anti-Human CD2 (CR-65192, Clone:TS1/8) or Mouse IgG2b Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Cardinal Red™ Anti-Human CD2 (CR-65192, Clone:TS1/8) or Mouse IgG2b Isotype Control. Cells were not fixed. Lymphocytes were gated.