

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD80

Numéro de catalogue: FITC-65217



## Informations de base

Numéro de catalogue:

FITC-65217

Taille:

100ug, 500 µg/ml

Hôte:

Mouse

Isotype:

IgG1, kappa

Numéro d'acquisition GenBank:

XM\_017597885

Identification du gène (NCBI):

25408

Nom complet:

Cd80 molecule

MW calculé

36 kDa

Méthode de purification:

Purification par affinité

CloneNo.:

3H5

Excitation/Emission maxima  
wavelengths:

495 nm / 524 nm

## Applications

Applications testées:

FC

Spécificité de l'espèce:

rat

## Informations générales

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

## Stockage

Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Tampon de stockage:

PBS with 0.09% sodium azide.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



1X10<sup>6</sup> unstimulated (dashed) or LPS-stimulated Wistar rat splenocytes were surface stained with 0.5 ug FITC Plus Anti-Rat CD80 (FITC-65217, Clone: 3H5) (red), or 0.5 ug Control Antibody. Cells were not fixed.