

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD357 (GITR)



Numéro de catalogue: **PE-65102**

Informations de base

Numéro de catalogue:

PE-65102

Taille:

100ug, 0.2 mg/ml

Hôte:

Rat

Isotype:

IgG2b

Numéro d'acquisition GenBank:

BC146517

Identification du gène (NCBI):

21936

Nom complet:

tumor necrosis factor receptor
superfamily, member 18

Méthode de purification:

Purification par affinité

CloneNo.:

DTA-1

Excitation/Emission maxima
wavelengths:

496 nm, 565 nm / 578 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

souris

Informations générales

Glucocorticoid-induced TNFR-related protein (GITR), also known as CD357 or TNFRSF18, is a member of the tumor necrosis factor receptor (TNF-R) superfamily. GITR is expressed constitutively at high levels in T regulatory cells (Treg cells) and plays a key role in dominant immunological self-tolerance maintained by CD25+CD4+ regulatory T cells (PMID: 11812990). It is expressed at low levels on resting responder T cells. The expression of GITR on T cells can be upregulated upon activation (PMID: 15770698). GITR is activated by GITR ligand (GITRL) which is mainly expressed on APC. GITR-GITRL interactions could co-stimulate both responder T-cell functions and the suppressive functions of Treg cells (PMID: 16868552).

Stockage

Stockage:

Stocker à une température comprise entre 2 °C et 8 °C. Éviter toute exposition à la lumière.

Tampon de stockage:

Tampon à base de phosphate avec azoture de sodium à 0,09 % et gélatine à 0,1 %, pH 7,2.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ mouse splenocytes were surface stained with 0.06 ug APC-Anti-Mouse CD4 (APC-65104, clone GK1.5), 0.125 ug FITC-Anti-Mouse CD25 (FITC-65137, clone PC61.5) and 0.06 ug PE-Anti-Mouse CD357 (GITR) (PE-65102, clone DTA-1). Cells were not fixed. CD4⁺ lymphocytes were gated for analysis of CD25 and CD357 (GITR) staining.