À des fins de recherche uniquement

Anticorps Monoclonal anti-CD5

Numéro de catalogue:PE-65152



Informations de base

Numéro de catalogue:

Numéro d'acquisition GenBank: BC027901

Méthode de purification:

PE-65152

Identification du gène (NCBI):

Purification par affinité

Taille:

CloneNo.: UCHT2

100 tests , 5 μ l/test Hôte: Mouse

Nom complet: CD5 molecule Excitation/Emission maxima wavelengths:

Isotype: IgG1, kappa MW calculé 495 aa, 55 kDa 496 nm, 565 nm / 578 nm

Applications

Applications testées:

FC

Spécificité de l'espèce:

Humain

Informations générales

CD5 is a type I transmembrane glycoprotein of the scavenger receptor cysteine-rich family (PMID: 12403363). CD5 is expressed on a majority of thymocytes, mature T cells, B cell subsets, and peripheral blood dendritic cells (PMID: 9858516; 6156984; 10379049; 1384337). CD5 may act as a receptor in regulating T-cell proliferation. It functions as a negative regulator of TCR signaling during thymocyte development (PMID: 7542801).

Stockage


Stockage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Tampon de stockage

PBS with 0.09% sodium azide and 0.5% BSA.

Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with APC Anti-Human CD19 and 5 ul PE Anti-Human CD5 (PE-65152, Clone:UCHT2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul PE Anti-Human CD5 (PE-65152, Clone:UCHT2) or Mouse IgG1 Isotype Control. Cells were not fixed. Lymphocytes were gated.