

For Research Use Only

PTER Polyclonal antibody

Catalog Number: 11191-1-AP



Basic Information

Catalog Number: 11191-1-AP	GenBank Accession Number: BC015092	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 200 µg/ml by Nanodrop and 200 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9317	Recommended Dilutions: WB 1:500-1:2000 IHC 1:20-1:200 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q96BW5	
Isotype: IgG	Full Name: phosphotriesterase related	
Immunogen Catalog Number: AG1675	Calculated MW: 39 kDa	
	Observed MW: 33 kDa, 39 kDa	

Applications

Tested Applications:

WB, IF, IHC, ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, mouse kidney tissue, mouse thymus tissue

IHC : human ovary tumor tissue,

IF/ICC : HEK-293 cells,

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human ovary tumor using 11191-1-AP (PTER antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (10% Formaldehyde) fixed HEK-293 cells using 11191-1-AP (PTER antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 11191-1-AP (PTER antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.