

For Research Use Only

CD147 Polyclonal antibody

Catalog Number: 11989-1-AP

Featured Product

36 Publications



Basic Information

Catalog Number: 11989-1-AP	GenBank Accession Number: BC009040	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 500 ug/ml by Nanodrop and 320 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 682	Recommended Dilutions: WB 1:5000-1:50000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:400-1:1600 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: P35613	
Isotype: IgG	Full Name: basigin (Ok blood group)	
Immunogen Catalog Number: AG2619	Calculated MW: 385 aa, 42 kDa	
	Observed MW: 35-58 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, IP, ELISA

Cited Applications:
WB, IHC, IF, IP, CoIP

Species Specificity:
human, mouse

Cited Species:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, human liver tissue, Jurkat cells, mouse heart tissue, mouse liver tissue, HepG2 cells

IP: HEK-293 cells,

IHC: human stomach cancer tissue, human liver cancer tissue, human lung cancer tissue

IF/ICC: A431 cells, HeLa cells, MCF-7 cells

Background Information

CD147, also known as Basigin or extracellular matrix metalloproteinase inducer (EMMPRIN), is a transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 7812975). The molecule is composed of an intracellular portion and an extracellular portion and a single transmembrane region. CD147 is expressed on a variety of cell types (e.g., hematopoietic, epithelial, and endothelial cells) and at varying levels (PMID: 32968061). Increased expression of CD147 occurs in many tumors. CD147 is a pleiotropic molecule that plays an important role in fetal, neuronal, lymphocyte and extracellular matrix development (PMID: 17945211). CD147 has been identified as a receptor essential for erythrocyte invasion by Plasmodium falciparum (PMID: 22080952). It has been reported that spike protein of SARS-CoV-2 binds to CD147 on host cells, thereby mediating the viral invasion (Wang, Ke, et al, BioRxiv, 2020). The CD147 gene, designated BSG for basigin, is located on chromosome 19p13.3. Four transcript variants encoding different isoforms have been found (PMID: 21536654). CD147 migrates on SDS-PAGE usually between 35 and 65 kDa, depending on the degree of glycosylation (PMID: 17945211).

Notable Publications

Author	Pubmed ID	Journal	Application
Linda Sooman	25263081	Acta Oncol	IHC
Lingling Gao	35003362	J Cancer	WB,IHC,CoIP
Tao Wang	28985237	PLoS Pathog	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11989-1-AP (CD147 antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.




WB result of CD147 antibody (11989-1-AP; 1:100000; room temperature for 1.5 hours) with wild-type and CD147 knockout Jurkat cells.




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 11989-1-AP (CD147 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 11989-1-AP (CD147 antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CD147 (IP:11989-1-AP, 4ug; Detection:11989-1-AP 1:10000) with HEK-293 cells lysate 1200 ug.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CD147 antibody (11989-1-AP) at dilution of 1:200 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).