For Research Use Only

SNAI2/SLUG Polyclonal antibody

Catalog Number:12129-1-AP

139 Publications



Basic Information

Catalog Number: 12129-1-AP

GenBank Accession Number:

BC015895

Purification Method: Antigen affinity purification

WB 1:5000-1:20000

GeneID (NCBI):

Recommended Dilutions:

150ul , Concentration: 650 µg/ml by Nanodrop;

UNIPROT ID:

043623 Full Name:

Rabbit

snail homolog 2 (Drosophila)

Isotype: IgG

Calculated MW:

Immunogen Catalog Number: AG2773

268 aa, 30 kDa Observed MW:

30 kDa

Applications

Tested Applications:

WB, FC, ELISA

Cited Applications:

WB, IF, IHC, ChIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, zebrafish

Positive Controls:

WB: mouse testis tissue, mouse brain tissue, rat heart

tissue, mouse spleen tissue

Background Information

 ${\sf SNA12}\ belongs\ to\ the\ Snail\ family\ of\ zinc\ finger\ transcription\ factors,\ which\ shares\ an\ evolution arily\ conserved\ role$ in mesoderm formation in vertebrates and invertebrates. It tiggers epithelial-mesenchymal transitions and has a crucial role in developmental processes. Also it is a transcriptional reperssor in mediating RAF-induced expression of TJ protein, occludin(OCLN) and subsequent oncogenic transformation of epithelial cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Yong Chen	36224027	Front Biosci (Landmark Ed)	WB
Xinxin Zhang	32980450	Gene	WB
Rongkun Li	34568024	Front Oncol	WB

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



1X10^6 MCF-7 cells were intracellularly stained with 0.2 ug Anti-Human SNAI2 (12129-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Various lysates were subjected to SDS PAGE followed by western blot with 12129-1-AP (SNAI2/SLUG antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.