For Research Use Only

Aggrecan Polyclonal antibody

Catalog Number:13880-1-AP 340 Publications



Basic Information

Catalog Number: 13880-1-AP GenBank Accession Number:

BC036445

Size: GenelD (NCBI):

150ul , Concentration: 500 ug/ml by 176 Nanodrop:

UNIPROT ID: P16112

Rabbit Full Name:
Isotype: aggrecan
IgG Calculated MW:
Immunogen Catalog Number: 250 kDa

AG4985

Purification Method: Antigen affinity purification Recommended Dilutions:

IHC: 1:200-1:800 FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Applications

Tested Applications: IHC, FC (Intra), ELISA

Cited Applications:

IHC, IF

Species Specificity: human, mouse, rat

Cited Species:

human, mouse, rat, pig, rabbit, canine, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

IHC : human lung tissue, human cartilage tissue, human lung cancer tissue, mouse lung tissue, rat brain

FC (Intra): ATDC-5 cells,

Background Information

Aggrecan (ACAN), also named as CSPCP, AGC1 and MSK16, belongs to the aggrecan/versican proteoglycan family (also including versican, brevican and neurocan). It is a major component of extracellular matrix of cartilaginous tissues. A major function of aggrecan is to resist compression in cartilage. It binds avidly to hyaluronic acid via an N-terminal globular region.

Notable Publications

Author	Pubmed ID	Journal	Application
Yangfei Yi	36178080	Orthop Surg	IHC
Sa Cha	34638972	Int J Mol Sci	IHC
Lindsay N Collins	31571216	J Comp Neurol	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 13880-1-AP (Aggrecan antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 13880-1-AP (Aggrecan antibody at dilution of 1:400 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 13880-1-AP (Aggrecan antibody at dilution of 1:400 (under 40x lens).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 13880-1-AP (Aggrecan antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 13880-1-AP (Aggrecan antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 13880-1-AP (Aggrecan antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 13880-1-AP (Aggrecan antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 ATDC-5 cells were intracellularly stained with 0.8 ug Anti-Human Aggrecan (13880-1-AP) and CoraLite 0.488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.