

For Research Use Only

# CD80 Polyclonal antibody

Catalog Number: 14292-1-AP **33 Publications**



## Basic Information

<b>Catalog Number:</b> 14292-1-AP	<b>GenBank Accession Number:</b> BC042665	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 700 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 941	<b>Recommended Dilutions:</b> WB 1:500-1:2000
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P33681	
<b>Isotype:</b> IgG	<b>Full Name:</b> CD80 molecule	
<b>Immunogen Catalog Number:</b> AG5615	<b>Calculated MW:</b> 33 kDa	
	<b>Observed MW:</b> 60 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : Raji cells, RAW 264.7 cells, human peripheral blood leukocyte, Ramos cells
<b>Cited Applications:</b> WB	
<b>Species Specificity:</b> human, mouse	
<b>Cited Species:</b> human, mouse, rat	

## Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

## Notable Publications

Author	Pubmed ID	Journal	Application
Haoran Wang	35087966	Bioact Mater	FC
Wen Song	36260348	J Med Chem	FC
Jiayu Zhang	35530972	Int J Nanomedicine	

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
Aliquoting is unnecessary for -20°C storage

**\*\*\* 20ul sizes contain 0.1% BSA**

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 14292-1-AP (CD80/B7-1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.