For Research Use Only

MAVS; VISA Polyclonal antibody

Catalog Number:14341-1-AP

Featured Product

87 Publications




Basic Information	Catalog Number: 14341-1-AP	GenBank Accession Number: BC044952		Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 800 ug/ml by	GenelD (NCBI): 57506		Recommended Dilutions: WB: 1:2000-1:16000
	Nanodrop;	UNIPROT ID:		IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	Source: Rabbit	Q7Z434		IHC: 1:250-1:1000
		Full Name:	al cignaling	IF/ICC: 1:50-1:500
	Isotype: IgG	mitochondrial antiviral signaling protein		
	Immunogen Catalog Number: AG5655	Calculated MW: 57 kDa		
		Observed MW: 50-55 kDa, 70-75 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA Cited Applications:		WB : A431 cells, HeLa cells, HuH-7 cells, Ju HepG2 cells	
	WB, IHC, IF, IP		IP : HEK-293 c	ells,
	Species Specificity:	H('huma		preast cancer tissue, human skin tissue
	human		IF/ICC : HeLa	cells.
	Cited Species: human, pig, monkey, duck			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI), it is widely involved a epithelial cells and he ctions. MAVS is able to i RAF6, TMEM173/MITA, I	and required for patocytes, conta nteract with va FIT3 and etc. It	r innate immune defense against virus ains CARD and transmembrane domair rious cellular proteins including can undergoe phosphorylation on
	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, wh the predicated 57 kDa.), it is widely involved a epithelial cells and hej ttions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause	and required for patocytes, conta nteract with va FIT3 and etc. It a the molecular	r innate İmmune defense against virus ains CARD and transmembrane domair rious cellular proteins including can undergoe phosphorylation on weight migrate to about 70 kDa despi
	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, wh the predicated 57 kDa.), it is widely involved a epithelial cells and he tions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause bmed ID Journ	and required for patocytes, conta nteract with va FIT3 and etc. It e the molecular	r innate immune defense against virus ains CARD and transmembrane domain rious cellular proteins including can undergoe phosphorylation on weight migrate to about 70 kDa despi
Background Information	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, whi the predicated 57 kDa.), it is widely involved a epithelial cells and hej ttions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause bmed ID Journ 587973 BMC N	and required for patocytes, conta nteract with va FIT3 and etc. It e the molecular al	r innate immune defense against virus ains CARD and transmembrane domain rious cellular proteins including can undergoe phosphorylation on weight migrate to about 70 kDa despin Application WB
	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, which the predicated 57 kDa. Author Put Jiangang Zheng 34 Ya-Ling Yang 36), it is widely involved a epithelial cells and hej ttions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause bmed ID Journ 587973 BMC V 174668 Eur J	and required for patocytes, conta nteract with va FIT3 and etc. It e the molecular al /et Res Pharmacol	can undergoe phosphorylation on weight migrate to about 70 kDa despit Application WB WB
	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, which the predicated 57 kDa. Author Put Jiangang Zheng 34 Ya-Ling Yang 36), it is widely involved a epithelial cells and hej ttions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause bmed ID Journ 587973 BMC N	and required for patocytes, conta nteract with va FIT3 and etc. It e the molecular al /et Res Pharmacol	r innate immune defense against virus ains CARD and transmembrane domain rious cellular proteins including can undergoe phosphorylation on weight migrate to about 70 kDa despi Application WB
	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, which the predicated 57 kDa. Author Pu Jiangang Zheng 34 Ya-Ling Yang 36 Lei-Ke Zhang 27 Storage: Storage suffer: PBS with 0.02% sodium azide and 50), it is widely involved a epithelial cells and hej ttions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause bmed ID Journ 587973 BMC \ 174668 Eur J 605671 J Viro ter shipment.	and required for patocytes, conta nteract with va FIT3 and etc. It e the molecular al /et Res Pharmacol	r innate immune defense against virus ains CARD and transmembrane domain rious cellular proteins including can undergoe phosphorylation on weight migrate to about 70 kDa despi Application WB WB
Notable Publications	promoter stimulator protein 1 (IPS-1 MAVS, present in T cells, monocytes, which are essential for antiviral func DDX58/RIG-I, IFIH1/MDA5, TRAF2, TI multiple sites and ubiquitination, which the predicated 57 kDa. Author Put Jiangang Zheng 34 Ya-Ling Yang 36 Lei-Ke Zhang 27 Storage: Storage Store at -20°C. Stable for one year af Storage Buffer:), it is widely involved a epithelial cells and hej ttions. MAVS is able to i RAF6, TMEM173/MITA, I hich may together cause bmed ID Journ 587973 BMC \ 174668 Eur J 605671 J Viro ter shipment.	and required for patocytes, conta nteract with va FIT3 and etc. It e the molecular al /et Res Pharmacol	r innate immune defense against virus ains CARD and transmembrane domain rious cellular proteins including can undergoe phosphorylation on weight migrate to about 70 kDa despi Application WB WB

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA)


W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 14341-1-AP (MAVS; VISA antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 14341-1-AP (MAVS; VISA antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 14341-1-AP (MAVS; VISA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




IP result of anti-MAVS; VISA (IP:14341-1-AP, 3ug; Detection:14341-1-AP 1:1000) with HEK-293 cells lysate 1700ug.




Various lysates were subjected to SDS PAGE followed by western blot with 14341-1-AP (MAVS; VISA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of MAVS; VISA antibody (14341-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MAVS; VISA transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 14341-1-AP (MAVS; VISA antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MAVS; VISA antibody (14341-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).