For Research Use Only

IFIT3 Polyclonal antibody Catalog Number:15201-1-AP Featured Product

Featured Product







Basic Information	Catalog Number: 15201-1-AP	GenBank Accession Numbe BC004977	r: Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 600 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG7349	GeneID (NCBI): 3437 UNIPROT ID: 014879	Recommended Dilutions: WB 1:2000-1:12000 IHC 1:50-1:500	
		Full Name: IFIT3		
		Calculated MW: 56 kDa		
		Observed MW: 60 kDa		
Applications	Tested Applications: WB, IHC, ELISA	Positive Controls:		
	Cited Applications: WB, IHC, IF, ChIP	WB : MKN-45 cells, SGC-7901 cells IHC : human spleen tissue,		
	Species Specificity: human			
	Cited Species: human, mouse, pig, monkey			
	Note-IHC: suggested antigen (TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
	IFIT3 (IFN-induced protein with tetratricopeptide repeats 3), also known as IFIT-4, ISG-60 or RIG-G, is a 56 kDa IFN induced protein. IFIT3 has been identified to be associated with strong IFN antiviral activity. Upon RNA virus infection, IFIT3 is significantly induced, ectopic expression or knockdown of IFIT3 could, respectively, enhance or impair IFN regulatory factor 3 (IRF3)-mediated gene expression. IFIT3 thus may serve as an important modulator in innate immunity, revealing a new function of the IFIT family proteins.			
Background Information	induced protein. IFIT3 has been iden infection, IFIT3 is significantly induc impair IFN regulatory factor 3 (IRF3)	ed, ectopic expression or kno -mediated gene expression. I	ckdown of IFIT3 could, respectively, enhance or FIT3 thus may serve as an important modulator in	
	induced protein. IFIT3 has been iden infection, IFIT3 is significantly induc impair IFN regulatory factor 3 (IRF3) innate immunity, revealing a new fu	ed, ectopic expression or kno -mediated gene expression. I	ckdown of IFIT3 could, respectively, enhance or FIT3 thus may serve as an important modulator ir	
Background Information Notable Publications	induced protein. IFIT3 has been iden infection, IFIT3 is significantly induc impair IFN regulatory factor 3 (IRF3) innate immunity, revealing a new fu Author Pu	ed, ectopic expression or kno -mediated gene expression. I nction of the IFIT family prot	ckdown of IFIT3 could, respectively, enhance or FIT3 thus may serve as an important modulator in eins.	
	induced protein. IFIT3 has been iden infection, IFIT3 is significantly induc impair IFN regulatory factor 3 (IRF3) innate immunity, revealing a new fu Author Pu Zhangtao Jiang 27	ed, ectopic expression or kno -mediated gene expression. I nction of the IFIT family prot bmed ID Journal	ckdown of IFIT3 could, respectively, enhance or FIT3 thus may serve as an important modulator in eins. Application WB	
	induced protein. IFIT3 has been iden infection, IFIT3 is significantly induc impair IFN regulatory factor 3 (IRF3) innate immunity, revealing a new fu Author Pu Zhangtao Jiang 27 Jiaming Su 36	ed, ectopic expression or kno -mediated gene expression. I nction of the IFIT family prot bmed ID Journal 681138 J Virol	ckdown of IFIT3 could, respectively, enhance or FIT3 thus may serve as an important modulator i eins. Application WB	
	induced protein. IFIT3 has been iden infection, IFIT3 is significantly induc impair IFN regulatory factor 3 (IRF3) innate immunity, revealing a new fu Author Pu Zhangtao Jiang 27 Jiaming Su 36	ed, ectopic expression or kno -mediated gene expression. I Inction of the IFIT family prot bmed ID Journal 681138 J Virol 163413 J Med Virol 813773 J Immunol ter shipment. 9% glycerol, pH7.3	ckdown of IFIT3 could, respectively, enhance or FIT3 thus may serve as an important modulator i eins. Application WB WB	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 15201-1-AP (IFIT3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 15201-1-AP (IFIT3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 15201-1-AP (IFIT3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).