

For Research Use Only

CYP19A1 Polyclonal antibody

Catalog Number: 16554-1-AP

Featured Product

19 Publications



Basic Information

Catalog Number: 16554-1-AP	GenBank Accession Number: BC022896	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 650 ug/ml by Nanodrop;	GeneID (NCBI): 1588	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500 IF-P 1:50-1:500 IF/ICC 1:50-1:500
Source: Rabbit	UNIPROT ID: P11511	
Isotype: IgG	Full Name: cytochrome P450, family 19, subfamily A, polypeptide 1	
Immunogen Catalog Number: AG9765	Calculated MW: 20 kDa, 25 kDa, 58 kDa	
	Observed MW: 49-55 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, IF-P, ELISA

Cited Applications:
WB, IHC, IF

Species Specificity:
human, mouse, rat

Cited Species:
human, mouse, rat, sheep, goat, geese

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A2780 cells, human placenta tissue

IHC: human placenta tissue, rat brain tissue, human kidney tissue, human skeletal muscle tissue, human gliomas tissue, human testis tissue, human spleen tissue, human lung tissue, human ovary tissue

IF-P: human placenta tissue,

IF/ICC: HepG2 cells,

Background Information

CYP19A1(Cytochrome P450 19A1) is also named as ARO1(aromatase), CYAR, CYP19, estrogen synthase and belongs to the cytochrome P450 family. It is a terminal enzyme which transforms irreversibly androgens into estrogens and it is present in the endoplasmic reticulum of numerous tissues(PMID:16406261). The gene encodes a protein with the molecular weight between 46 kDa and 69 kDa(PMID:8129748) and the protein can be glycosylated, but the level of glycosylation does not seem to be essential for CYP19A1 activity(PMID:12606587). Brain aromatase may be neuroprotective by increasing the local estrogen levels in injured neurons so that genetic variation in the brain aromatase gene may modify the risk for AD(PMID:16767510).

Notable Publications

Author	Pubmed ID	Journal	Application
Huidan Wang	26358501	Sci Rep	WB
Ruiming Fan	32875818	Br Poult Sci	WB
Yan Peng	33109277	Biol Res	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A2780 cells were subjected to SDS PAGE followed by western blot with 16554-1-AP (CYP19A1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




human placenta tissue were subjected to SDS PAGE followed by western blot with 16554-1-AP (CYP19A1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 16554-1-AP (CYP19A1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 16554-1-AP (CYP19A1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human placenta tissue using CYP19A1 antibody (16554-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CYP19A1 antibody (16554-1-AP) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).