For Research Use Only

## ZC3HAV1 Polyclonal antibody

Catalog Number:16820-1-AP

Featured Product

33 Publications




Basic Information	Catalog Number: 16820-1-AP	GenBank Accession Number: BC025308	Purification Method: Antigen affinity purification	
	Size:	GeneID (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 650 ug/ml by	56829	WB 1:2000-1:12000	
	Nanodrop and 367 ug/ml by Bradford method using BSA as the standard;	UNIPROT ID: Q7Z2W4	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Source: Rabbit	Full Name: zinc finger CCCH-type, antivit	ral 1	
	Isotype: IgG	Calculated MW: 699aa,78 kDa; 902aa,101 kDa		
	Immunogen Catalog Number: AG10364	Observed MW: 100 kDa		
Applications	Tested Applications: WB, IP, ELISA	Positive Controls: WB : HEK-293 cells, HeLa cells IP : HeLa cells,		
	Cited Applications: WB, IHC, IF, IP, RIP			
	Species Specificity: human, mouse, rat			
	Cited Species: human, mouse, monkey			
Background Information	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innat	s a CCCH-type zinc finger prote e immunity to viral infection. A	a rat cDNA library due to its conferring ein, it may primarily function to inhibit viral lternative splicing occurs at this locus and tv forms with MV 101, 77, 67, 42 and 18 kDa.	
	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innate variants, each encoding distinct isofo	s a CCCH-type zinc finger prote e immunity to viral infection. A	ein, it may primarily function to inhibit viral Ilternative splicing occurs at this locus and tw	
	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innat variants, each encoding distinct isofo Author Put	s a CCCH-type zinc finger prote e immunity to viral infection. A rms. ZC3HAV1 exists some iso	ein, it may primarily function to inhibit viral lternative splicing occurs at this locus and tw forms with MV 101, 77, 67, 42 and 18 kDa. Application	
	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innativity variants, each encoding distinct isofor Author Put Yongxiang Zhao 330	s a CCCH-type zinc finger prote e immunity to viral infection. A rrms. ZC3HAV1 exists some iso pmed ID Journal	ein, it may primarily function to inhibit viral lternative splicing occurs at this locus and tw forms with MV 101, 77, 67, 42 and 18 kDa. Application	
Background Information	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innativity variants, each encoding distinct isofor Author Put Yongxiang Zhao 330	s a CCCH-type zinc finger prote e immunity to viral infection. A rms. ZC3HAV1 exists some iso pmed ID Journal D35816 Vet Microbiol	ein, it may primarily function to inhibit Iternative splicing occurs at this locus forms with MV 101, 77, 67, 42 and 18 kl Applicati WB	
Notable Publications	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innate variants, each encoding distinct isofor Author Put Yongxiang Zhao 330 Matthew A Takata 289	s a CCCH-type zinc finger prote e immunity to viral infection. A rms. ZC3HAV1 exists some iso pmed ID Journal D35816 Vet Microbiol	ein, it may primarily function to inhibit vira lternative splicing occurs at this locus and forms with MV 101, 77, 67, 42 and 18 kDa. Application WB WB	
	human, mouse, monkey The zinc-finger antiviral protein (ZAP resistance to retroviruses. ZC3HAV1 i gene expression and induce an innate variants, each encoding distinct isofo Author Put Yongxiang Zhao 330 Matthew A Takata 289 Evan P Booy 302	s a CCCH-type zinc finger prote e immunity to viral infection. A rms. ZC3HAV1 exists some iso omed ID Journal 035816 Vet Microbiol 053888 Nature 247708 Nucleic Acids er shipment.	ein, it may primarily function to inhibit viral llternative splicing occurs at this locus and to forms with MV 101, 77, 67, 42 and 18 kDa. Application WB WB	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16820-1-AP (ZC3HAV1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




WB result of ZC3HAV1 antibody (16820-1-AP; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ZC3HAV1 transfected HEK-293 cells.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16820-1-AP (ZC3HAV1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 16820-1-AP (ZC3HAV1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-ZC3HAV1 (IP:16820-1-AP, 4ug; Detection:16820-1-AP 1:2000) with HeLa cells lysate 1080ug.



Immunohistochemical analysis of paraffinembedded stomach cancer slide using 16820-1-AP (ZC 3HAV 1 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).